VACCINES DURING PREGNANCY

Timing and Notes About Protection

Receives primary benefit

Protection Notes for Mom

Protection Notes for Baby

TTTT

INFLUENZA

Timing

July - Aug. (if available): Those in 3rd trimeste

Sept. - spring: Pregnant individuals who did not previously get the vaccine

COVID-19

Timing

Anytime during pregnancy if not up to date on most recent version; updated version when available in the fall

\checkmark

At increased risk for severe infection

At increased risk for severe infection or

preterm labor

Вару

Receives protection from maternal antibodies until old enough to get vaccinated (6 months of age) Level of protection will vary based on

Level of protection will vary based on timing of maternal vaccination relative to influenza season and infant's date of birth.

Receives protection from maternal antibodies until old enough to get vaccinated (6 months of age)

Level of protection will vary based on relative timing of maternal vaccination and infant's date of birth.

TDAP

Timing

RSV

Between 27 and 36 weeks of gestation during each pregnancy Increased protection after vaccination, but not at increased risk for severe disease Protection against pertussis (whooping cough) in the first few months of life when baby is at greatest risk of hospitalization or death from pertussis

\checkmark

Protection during first RSV season when risk of severe disease is greatest

Maternal option has been shown to protect about 5 or 6 of every 10 infants. Infant option has been shown to protect about 7 or 8 of every 10 infants.

Children's Hospital of Philadelphia⁻ Vaccine Education Center

©2025 Children's Hospital of Philadelphia. 25302-01-25

Not at increased risk for severe RSV

Find out more:



vaccine.chop.edu/parents (Check "Parents PACK Newsletter" February 2025 issue)

Timing

Maternal option: RSV vaccine if 32 to 36 weeks of gestation occurs between Sept. and Jan. (based on RSV season; applies for most parts of the US)

OR

Infant option: Nirsevimab in first week of life if born during RSV season or in the fall (usually Oct.) if not yet 8 months of age

In most cases only one of these options is needed.